

# The Kernel Trick

Robert M. Haralick

Computer Science, Graduate Center  
City University of New York



# SVM Classification

$\langle (x_1, c_1), \dots, (x_Z, c_Z) \rangle$  is the training data  
 $c_1, \dots, c_Z \in \{-1, 1\}$  specifies the corresponding classes  
 $\alpha_1, \dots, \alpha_Z$  are the Lagrange multipliers

Assign  $x$  to class +1 when  $w'x > \theta$  Else assign to class -1

$$\begin{aligned}w'x &= \left( \sum_{z \in S} \alpha_z c_z x_z \right)' x \\ &= \sum_{z \in S} \alpha_z c_z (x_z' x)\end{aligned}$$

# Dual Form

$$\text{Minimize } L(\alpha_1, \dots, \alpha_Z) = \sum_{z=1}^Z \alpha_z - \frac{1}{2} \sum_{i=1}^Z \sum_{j=1}^Z \alpha_i \alpha_j \mathbf{c}_i \mathbf{c}_j (\mathbf{x}'_i \mathbf{x}_j)$$

subject to

$$\alpha_z \geq 0, z = 1, \dots, Z$$

$$\sum_{z=1}^Z \alpha_z \mathbf{c}_z = \mathbf{0}$$

# Making Non-Linear Boundaries

Example: Change  $x^{2 \times 1} = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$  To  $\phi(x)^{5 \times 1} = \begin{pmatrix} x_1^2 \\ x_2^2 \\ x_1 x_2 \\ x_1 \\ x_2 \end{pmatrix}$

$$\text{Minimize } L(\alpha_1, \dots, \alpha_Z) = \sum_{z=1}^Z \alpha_z - \frac{1}{2} \sum_{i=1}^Z \sum_{j=1}^Z \alpha_i \alpha_j c_i c_j (\phi(x_i)' \phi(x_j))$$

subject to

$$\alpha_z \geq 0, z = 1, \dots, Z$$
$$\sum_{z=1}^Z \alpha_z c_z = 0$$

$$w = \sum_{z \in S} \alpha_z c_z \phi(x_z)$$

# Making Non-Linear Boundaries

Assign  $x$  to class +1 when

$$w' \phi(x) > \theta$$
$$w_1 x_1^2 + w_2 x_2^2 + w_3 x_1 x_2 + w_4 x_1 + w_5 x_2 > \theta$$

- The linear problem in the 5-dimensional space is the *same* as the quadratic problem in the 2-dimensional space.
- The kernel trick avoids the explicit mapping that is needed to get linear learning algorithms to learn a nonlinear function or decision boundary.

# Making Non-Linear Boundaries

$$\text{Minimize } L(\alpha_1, \dots, \alpha_Z) = \sum_{z=1}^Z \alpha_z - \frac{1}{2} \sum_{i=1}^Z \sum_{j=1}^Z \alpha_i \alpha_j \mathbf{c}_i \mathbf{c}_j (\phi(\mathbf{x}_i)' \phi(\mathbf{x}_j))$$

$$\text{Set } K(\mathbf{x}, \mathbf{y}) = \phi(\mathbf{x})' \phi(\mathbf{y})$$

$$\text{Minimize } L(\alpha_1, \dots, \alpha_Z) = \sum_{z=1}^Z \alpha_z - \frac{1}{2} \sum_{i=1}^Z \sum_{j=1}^Z \alpha_i \alpha_j \mathbf{c}_i \mathbf{c}_j K(\mathbf{x}_i, \mathbf{x}_j)$$

$$\mathbf{w}' \phi(\mathbf{x}) = \sum_{z \in S} \alpha_z \mathbf{c}_z \phi(\mathbf{x}_z)' \phi(\mathbf{x})$$

$$= \sum_{z \in S} \alpha_z \mathbf{c}_z K(\mathbf{x}_z, \mathbf{x})$$

# The Kernel Trick

What happens when  $K(x, y) = (x'y)^2$ ?

Example:

$$x = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$$

$$y = \begin{pmatrix} y_1 \\ y_2 \end{pmatrix}$$

$$x'y = x_1y_1 + x_2y_2$$

$$(x'y)^2 = x_1^2y_1^2 + 2x_1y_1x_2y_2 + x_2^2y_2^2$$

$$= \begin{pmatrix} x_1^2 \\ \sqrt{2}x_1x_2 \\ x_2^2 \end{pmatrix}' \begin{pmatrix} y_1^2 \\ \sqrt{2}y_1y_2 \\ y_2^2 \end{pmatrix}$$

$$= \phi(x)'\phi(y)$$

The components of the weight vector can compensate for coefficients that are not 1.



# Two-Class Data Set




Figure: Raw Data in 2D is not separable with hyperplane.

# Two-Class Data Set Transformed




Figure: Transformed:  $(x_1, x_2) \mapsto (x_1^2, \sqrt{2}x_1x_2, x_2^2)$

# Brute Force: Quadratic Case

If  $x$  has  $N$  components, to make it quadratic requires an additional  $N(N - 1)/2$  components

- $x^{N \times 1}$
- $\phi(x)^{(N+N*(N-1)/2) \times 1}$
- $w^{(N+N*(N-1)/2) \times 1} = \sum_{z \in S} \alpha_z c_z \phi(x_z)$
- $w' \phi(x) = \sum_{z \in S} \alpha_z c_z \phi(x_z)' \phi(x)$
- Takes  $N + N * (N - 1)/2$  operations for  $x \mapsto \phi(x)$
- Takes  $N + N * (N - 1)/2$  operations for  $w' \phi(x)$

# The Kernel Trick

- $x^{N \times 1}$
- $x'y$  takes  $N$  operations
- $(x'y)^2$  takes  $N + 1$  operations
- $\sum_{z \in S} \alpha_z c_z K(x_z, x)$
- $K(x_z, x) = (x'_z x)^2$  takes  $N + 1$  operations
- Compute  $w'x = |S|(N + 1)$  operations
- Brute Force  $N + N * (N - 1)/2$  operations

# The Kernel Trick

What happens when  $K(x, y) = (x'y)^3$ ?

Example:

$$\begin{aligned}x'y &= x_1y_1 + x_2y_2 \\(x'y)^3 &= (x_1y_1)^3 + 3(x_1y_1)^2x_2y_2 + 3(x_1y_1)(x_2y_2)^2 + (x_2y_2)^3 \\&= \begin{pmatrix} x_1^3 \\ \sqrt{3}x_1^2x_2 \\ \sqrt{3}x_1x_2^2 \\ x_2^3 \end{pmatrix}' \begin{pmatrix} y_1^3 \\ \sqrt{3}y_1^2y_2 \\ \sqrt{3}y_1y_2^2 \\ y_2^3 \end{pmatrix}\end{aligned}$$

# The Kernel Trick

What kind of functions  $K(x, y)$  can be represented in the form

$$K(x, y) = \phi(x)' \phi(y)$$

## Definition

A **Kernel** is a function  $K : \mathbb{R}^M \times \mathbb{R}^M \rightarrow \mathbb{R}$ , such that for all  $x, y \in \mathbb{R}^M$ , there is a feature function  $\phi : \mathbb{R}^M \rightarrow \mathbb{R}^N$  such that

$$K(x, y) = \phi(x)' \phi(y)$$

## Proposition

*Kernels are symmetric functions.*

Let  $x = (x_1, \dots, x_M)'$  and  $y = (y_1, \dots, y_M)'$

$$\begin{aligned}K(x, y) &= (x'y)^2 \\&= \left( \sum_{m=1}^M x_m y_m \right)^2 \\&= \sum_{i=1}^M x_i y_i \sum_{j=1}^M x_j y_j \\&= \sum_{i=1}^M \sum_{j=1}^M x_i y_i x_j y_j \\&= \sum_{i=1}^M \sum_{j=1}^M (x_i x_j) (y_i y_j) \\&= \left( (x_i x_j) \Big|_{(i,j)=(1,1)}^{(M,M)} \right)' \left( (y_i y_j) \Big|_{(i,j)=(1,1)}^{(M,M)} \right) \\ \phi(x) &= \left( (x_i x_j) \Big|_{(i,j)=(1,1)}^{(M,M)} \right)\end{aligned}$$



Let  $x = (x_1, \dots, x_M)'$  and  $y = (y_1, \dots, y_M)'$

$$\begin{aligned}K(x, y) &= (x'y + 1)^2 \\&= \left( \sum_{i=1}^M x_i y_i + 1 \right) \left( \sum_{j=1}^M x_j y_j + 1 \right) \\&= \sum_{i=1}^M \sum_{j=1}^M x_i x_j y_i y_j + 2 \sum_{i=1}^M x_i y_i + 1 \\&= \sum_{i=1}^M \sum_{j=1}^M (x_i x_j)(y_i y_j) + \sum_{i=1}^M (\sqrt{2}x_i)(\sqrt{2}y_i) + (1)(1)\end{aligned}$$

# Common Kernels

$$x, y \in \mathbb{R}^N$$

Polynomial	$K(x, y) = (x'y + 1)^q$	$N + 1 + 2$ (log and exponentiate)
Radial	$K(x, y) = \exp\left(-\frac{\ x-y\ ^2}{\sigma^2}\right)$	$N + 1 + 1$ (exponentiate)
Sigmoid	$K(x, y) = \tanh(\beta x'y + \gamma)$	$N + 1 + 1$ (hyperbolic tangent)

$K(x, y)$  measures the similarity between  $x$  and  $y$

# Kernels and Feature Spaces

Given a function  $K : \mathbb{R}^M \times \mathbb{R}^M \rightarrow \mathbb{R}$ , how can we tell that there exists a feature function  $\phi$  such that

$$K(x, y) = \phi(x)' \phi(y)$$

thereby making  $K$  a kernel?

# Mercer's Kernel Characterization Theorem

Let  $K : \mathbb{R}^M \times \mathbb{R}^M \rightarrow \mathbb{R}$  be a symmetric function. There exists  $\phi : \mathbb{R}^M \rightarrow \mathbb{R}^N$  such that

$$K(x, y) = \phi(x)' \phi(y)$$

if and only if

$$\int_{x \in \mathbb{R}^M} \int_{y \in \mathbb{R}^M} K(x, y) g(x) g(y) dx dy \geq 0$$

for every function  $g : \mathbb{R}^M \rightarrow \mathbb{R}$  satisfying

$$\int_{x \in \mathbb{R}^M} g(x)^2 dx < \infty$$

# Mercer's Kernel Characterization Theorem

Let  $K : \mathbb{R}^M \times \mathbb{R}^M \rightarrow \mathbb{R}$  be a symmetric function. There exists  $\phi : \mathbb{R}^M \rightarrow \mathbb{R}^N$  such that

$$K(x, y) = \phi(x)' \phi(y)$$

if and only if for every  $L$  and every set  $\{x_1, \dots, x_L\}$  of  $L$  points from  $\mathbb{R}^M$ , the matrix  $\mathbb{K}$ ,

$$\mathbb{K} = \begin{pmatrix} K(x_1, x_1) & K(x_1, x_2) & \dots & K(x_1, x_L) \\ K(x_2, x_1) & K(x_2, x_2) & \dots & K(x_2, x_L) \\ & \vdots & \vdots & \\ K(x_L, x_1) & K(x_L, x_2) & \dots & K(x_L, x_L) \end{pmatrix}$$

is positive semidefinite.

# Symmetric Positive Semi-definite

## Definition

A symmetric matrix  $B^{N \times N}$  is **Positive Semi-definite** if and only if for every  $x \in \mathbb{R}^N$ ,

$$x' B x \geq 0$$

## Theorem

*A symmetric matrix  $B$  is positive semi-definite if and only if all its eigenvalues are non-negative.*

## Theorem

*If  $B$  is a symmetric positive semi-definite matrix, then there exists a matrix  $A$  satisfying  $B = A' A$ .*

# Positive Definite Kernel

## Definition

A symmetric function  $K : \mathbb{R}^N \times \mathbb{R}^N \rightarrow \mathbb{R}$  is **Positive Definite** (pd) if and only if for any  $x_1, \dots, x_L \in \mathbb{R}^N$  and any  $c_1, \dots, c_L \in \mathbb{R}$

$$\sum_{i=1}^L \sum_{j=1}^L c_i K(x_i, x_j) c_j \geq 0$$

An alternative way of writing  $\sum_{i=1}^L \sum_{j=1}^L c_i K(x_i, x_j) c_j$  is  $c' \mathbb{K} c$  where  $c' = (c_1, \dots, c_L)$  and  $\mathbb{K}_{ij} = K(x_i, x_j)$

# A Positive Definite Kernel

## Proposition

Let  $K : \mathbb{R}^N \times \mathbb{R}^N \rightarrow \mathbb{R}$  be defined by  $K(x, y) = x'y$ . Then  $K$  is a positive definite kernel.

## Proof.

Let  $x_1, \dots, x_K \in \mathbb{R}^N$ . Define

$$A = \begin{pmatrix} x_1' \\ x_2' \\ \vdots \\ x_K' \end{pmatrix}$$

$$K = AA'$$

Let  $c \in \mathbb{R}^K$ .

$$\begin{aligned} c'Kc &= c'AA'c \\ &= (A'c)'(A'c) \geq 0 \end{aligned}$$



# Positive Definite Kernel Properties

## Proposition

Suppose  $K_i : \mathbb{R}^N \times \mathbb{R}^N \rightarrow \mathbb{R}, i = 1, 2, \dots$ , are positive definite (pd) kernels then

- $K(x, y) = \sum_{j=1}^J a_j K_j(x, y)$  is a pd kernel when  $a_j \geq 0$
- $K(x, y) = \prod_{j=1}^J K_j(x, y)$  is a pd kernel
- $K(x, y) = \lim_{j \rightarrow \infty} K_j(x, y)$  is a pd kernel
- $K(x, y) = q(K_1(x, y))$  is a pd kernel where  $q$  is any polynomial with non-negative coefficients
- $K(x, y) = f(x)K_1(x, y)f(y)$  is a pd kernel for any  $f : \mathbb{R}^N \rightarrow \mathbb{R}$
- $K(x, y) = f(x)f(y)$  is a pd kernel for any  $f : \mathbb{R}^N \rightarrow \mathbb{R}$
- $K(x, y) = K_1(\phi(x), \phi(y))$
- $K(x, y) = x' B z$ , where  $B$  is symmetric positive semidefinite matrix

# Products of pd kernels are pd kernels

## Proposition

Let  $\psi_1, \psi_2 : \mathbb{R}^N \times \mathbb{R}^N \rightarrow \mathbb{R}$  be positive definite kernels. Define  $\psi : \mathbb{R}^N \times \mathbb{R}^N \rightarrow \mathbb{R}$  by  $\psi(x, y) = \psi_1(x, y)\psi_2(x, y)$ . Then  $\psi$  is a positive definite kernel.

## Proof.

Let  $x_1, \dots, x_M \in \mathbb{R}^N$  and let  $c_1, \dots, c_M \in \mathbb{R}$ . Then

$$\sum_{i=1}^M \sum_{j=1}^M c_i \psi(x_i, x_j) c_j = \sum_{i=1}^M \sum_{j=1}^M c_i \psi_1(x_i, x_j) \psi_2(x_i, x_j) c_j$$

Since  $\psi_1$  and  $\psi_2$  are both positive definite, there exist matrices  $A^{M \times M}$  and  $B^{M \times M}$  such that

$$\begin{aligned} \psi_1(x_i, x_j) &= (A' A)_{ij} = \sum_{k=1}^M a_{ki} a_{kj} \\ \psi_2(x_i, x_j) &= (B' B)_{ij} = \sum_{k=1}^K b_{ki} b_{kj} \end{aligned}$$

# Products of pd kernels are pd kernels

Proof.

Hence,  $\psi_1(x_i, x_j) = \sum_{k=1}^M a_{ki} a_{kj}$  and  $\psi_2(x_i, x_j) = \sum_{l=1}^M b_{li} b_{lj}$ . Therefore,

$$\begin{aligned} \sum_{i=1}^M \sum_{j=1}^M c_i \psi_1(x_i, x_j) \psi_2(x_i, x_j) c_j &= \sum_{i=1}^M \sum_{j=1}^M c_i \sum_{k=1}^M a_{ki} a_{kj} \sum_{l=1}^M b_{li} b_{lj} \\ &= \sum_{k=1}^M \sum_{l=1}^M \left( \sum_{i=1}^M c_i a_{ki} b_{li} \right) \left( \sum_{j=1}^M c_j a_{kj} b_{lj} \right) \\ &= \sum_{k=1}^M \sum_{l=1}^M \left( \sum_{i=1}^M c_i a_{ki} b_{li} \right)^2 \geq 0 \end{aligned}$$

□

# Exponential Kernel

## Proposition

Define  $q_j(x) = \sum_{n=0}^j \frac{x^n}{n!}$ . Suppose  $K : \mathbb{R}^N \times \mathbb{R}^N \rightarrow \mathbb{R}$  is a positive definite kernel, then,

$$\exp(K(x, y)) = \lim_{j \rightarrow \infty} q_j(K(x, y))$$

is a positive definite kernel.

## Proof.

$q_j(x)$  is a polynomial with all positive coefficients. Since  $K$  is a positive definite kernel,  $q_j(K(x, y))$  is a positive definite kernel. Limits of positive definite kernels are positive definite kernels. Therefore  $\exp(K(x, y))$  is a positive definite kernel.  $\square$

# Positive Definite Kernel Property

## Proposition

Let  $\psi : \mathbb{R}^N \times \mathbb{R}^N \rightarrow \mathbb{R}$  be a symmetric positive definite kernel. Let  $f : \mathbb{R}^N \rightarrow \mathbb{R}$  be any function. Then  $K(x, y) = f(x)\psi(x, y)f(y)$  is a symmetric positive definite kernel.

## Proof.

The symmetry is immediate because  $\psi$  is symmetric and multiplication is commutative. Let  $x_1, \dots, x_M \in \mathbb{R}^N$  and  $c_1, \dots, c_M \in \mathbb{R}$ . Then

$$\begin{aligned} \sum_{i=1}^M \sum_{j=1}^M c_i K(x_i, x_j) c_j &= \sum_{i=1}^M \sum_{j=1}^M c_i f(x_i) \psi(x_i, x_j) f(x_j) c_j \\ &= \sum_{i=1}^M \sum_{j=1}^M (c_i f(x_i)) \psi(x_i, x_j) (f(x_j) c_j) \end{aligned}$$

Let  $d_i = c_i f(x_i)$ . Then, since  $\psi$  is positive definite,

$$\sum_{i=1}^M \sum_{j=1}^M c_i K(x_i, x_j) c_j = \sum_{i=1}^M \sum_{j=1}^M d_i \psi(x_i, x_j) d_j \geq 0$$

# Positive Definite Kernel

## Proposition

Let  $K : \mathbb{R}^N \times \mathbb{R}^N \rightarrow \mathbb{R}$  be a positive definite kernel. Then  $K(x, x)K(y, y) \geq K(x, y)^2$ .

## Proof.

Let  $x, y \in \mathbb{R}^N$ . Then the matrix  $\begin{pmatrix} K(x, x) & K(x, y) \\ K(y, x) & K(y, y) \end{pmatrix}$  is positive semidefinite. Hence the determinant must be non-negative so that  $K(x, x)K(y, y) - K(x, y)^2 \geq 0$ . Therefore  $K(x, x)K(y, y) \geq K(x, y)^2$ . □

# Example

## Proposition

Let  $K : \mathbb{R}^N \times \mathbb{R}^N \rightarrow \mathbb{R}$  be defined by

$$K(x, y) = \exp\left(-\frac{(x-y)'(x-y)}{\sigma^2}\right)$$

Then  $K$  is a positive definite kernel.

## Proof.

$x'y$  is a pd kernel.

$\frac{2x'y}{\sigma^2}$  is a pd kernel.

$\exp\left(\frac{2x'y}{\sigma^2}\right)$  is a pd kernel.

$\exp\left(-\frac{x'x}{\sigma^2}\right) \exp\left(\frac{2x'y}{\sigma^2}\right) \exp\left(-\frac{y'y}{\sigma^2}\right)$  is a pd kernel.

$\exp\left(-\frac{(x-y)'(x-y)}{\sigma^2}\right)$  is a pd kernel. □

# Positive Definite Kernel Properties

## Proposition

Let  $K : \mathbb{R}^N \times \mathbb{R}^N \rightarrow \mathbb{R}$  be a positive definite kernel. Then

$$K_1(x, y) = \frac{K(x, y)}{\sqrt{K(x, x)K(y, y)}}$$

is a positive definite kernel.

## Proof.

$$\begin{aligned} K_1(x, y) &= \frac{K(x, y)}{\sqrt{K(x, x)K(y, y)}} \\ &= \frac{1}{\sqrt{K(x, x)}} K(x, y) \frac{1}{\sqrt{K(y, y)}} \end{aligned}$$





# Positive Definite Kernel

## Proposition

*Let  $h : \mathbb{R}^N \rightarrow \mathbb{R}$  be any function satisfying  $h(x) \geq 0$  with minimum at 0. Then,*

$$K(x, y) = h(x + y) - h(x - y)$$

*is a positive definite kernel.*

# Symmetric Negative Definite

## Definition

A symmetric function  $\psi : \mathbb{R}^N \times \mathbb{R}^N \rightarrow \mathbb{R}$  is **Negative Definite** if and only if for every  $x_1, \dots, x_M \in \mathbb{R}^N$  and  $c_1, \dots, c_M \in \mathbb{R}$  satisfying  $\sum_{m=1}^M c_m = 0$

$$\sum_{i=1}^M \sum_{j=1}^M c_i \psi(x_i, x_j) c_j \leq 0$$

## Proposition

Let  $\Psi^{M \times M}$  be defined by  $\Psi_{ij} = \psi(x_i, x_j)$ . If  $\Psi$  is negative semi-definite Then  $\psi$  is negative definite.

# Properties of Kernels

## Proposition

*Let  $K : \mathbb{R}^N \times \mathbb{R}^N \rightarrow \mathbb{R}$  be a symmetric function. If  $K$  is a positive definite kernel then  $-K$  is a negative definite kernel.*

## Proof.

*Let  $x_1, \dots, x_K \in \mathbb{R}^N$  and  $c \in \mathbb{R}^K$ . Suppose  $K$  is a positive definite kernel. then  $c'Kc \geq 0$ . Then  $c'(-K)c = -c'Kc \leq 0$ . Since this is true for all  $c$ , it is certainly true for  $c$  satisfying  $1'c = 0$  and this implies that  $-K$  is a negative definite kernel.  $\square$*

# A Negative Definite Kernel

## Proposition

Let  $\psi : \mathbb{R}^N \times \mathbb{R}^N$  be defined by  $\psi(x, y) = (x - y)'(x - y)$ . Then  $\psi$  is a negative definite kernel.

## Proof.

Let  $x_1, \dots, x_K \in \mathbb{R}^N$  and  $c_1, \dots, c_K \in \mathbb{R}$  satisfying  $\sum_{k=1}^K c_k = 0$ . Then,

$$\begin{aligned} \sum_{i=1}^K \sum_{j=1}^K c_i (x_i - x_j)'(x_i - x_j) c_j &= \sum_{i=1}^K c_i x_i' x_i \sum_{j=1}^K c_j - 2 \sum_{i=1}^K \sum_{j=1}^K c_i x_i' x_j c_j + \\ &\quad \sum_{i=1}^K c_i \sum_{j=1}^K x_j' x_j c_j \\ &= -2 \sum_{i=1}^K \sum_{j=1}^K c_i x_i' x_j c_j \leq 0 \end{aligned}$$



# Negative Definite Kernel Properties

## Proposition

Let  $K_m : \mathbb{R}^N \times \mathbb{R}^N \rightarrow \mathbb{R}$ ,  $m = 1, 2, \dots$ , be symmetric and negative definite. Then  $J_1, J_2, J_3, J_4$  and  $J_5$  defined by

$$J_1(x, y) = \sum_{m=1}^M a_m K_m(x, y), \quad a_m \geq 0$$

$$J_2(x, y) = \lim_{m \rightarrow \infty} K_m(x, y)$$

$$J_3(x, y) = \log(1 + K(x, y))$$

$$J_4(x, y) = K(x, y)^\alpha, \quad 0 < \alpha < 1$$

$$J_5(x, y) = f(x) + f(y), \quad \text{for any } f : \mathbb{R}^N \rightarrow \mathbb{R}$$

are negative definite kernels.

# Negative Definite Kernel

## Proposition

Let  $\psi : \mathbb{R}^N \times \mathbb{R}^N \rightarrow \mathbb{R}$  be a symmetric negative definite kernel. Then  $\psi(x, x) + \psi(y, y) \leq 2\psi(x, y)$

## Proof.

Let  $x, y \in \mathbb{R}^N$  and  $c_1, c_2 \in \mathbb{R}$  satisfying  $c_1 + c_2 = 0$ . Since  $\psi$  is negative definite,

$$\begin{aligned} 0 &\geq c_1\psi(x, x)c_1 + c_1\psi(x, y)c_2 + c_2\psi(y, x)c_1 + c_2\psi(y, y)c_2 \\ &\geq c_1\psi(x, x)c_1 + c_1\psi(x, y)(-c_1) + (-c_1)\psi(y, x)c_1 + (-c_1)\psi(y, y)(-c_1) \\ &\geq \psi(x, x) - 2\psi(x, y) + \psi(y, y) \\ 2\psi(x, y) &\geq \psi(x, x) + \psi(y, y) \end{aligned}$$



# Positive and Negative Definite Symmetric Functions

## Theorem

*(Shoenberg, 1938)*

*Let  $\psi : \mathbb{R}^N \times \mathbb{R}^N \rightarrow \mathbb{R}$  be symmetric. Then  $\psi(x, y)$  is negative definite if and only if  $\exp(-t\psi(x, y))$  is positive definite for all  $t > 0$ .*

## Proposition

*Let  $K : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$  be a positive definite kernel satisfying  $K(x, y) \geq 0$ ,  $x, y \in \mathbb{R}^n$ . Then*

$$-\log K(x, y)$$

*is negative definite if and only if*

$$K(x, y)^t$$

*is positive definite for all  $t > 0$*

# Positive and Negative Definite Symmetric Functions

## Proposition

Let  $\psi : \mathbb{R}^N \times \mathbb{R}^N \rightarrow \mathbb{R}$  be a symmetric function. Let  $z \in \mathbb{R}^N$ . Define

$$\phi(x, y) = -\psi(x, y) + \psi(x, z) + \psi(z, y) - \psi(z, z)$$

Then  $\psi$  is negative definite if and only if  $\phi$  is positive definite.

## Proof.

Suppose  $\psi$  is negative definite. Let  $x_k \in \mathbb{R}^N$ ,  $k = 0, \dots, K$  and  $c_k \in \mathbb{R}$ ,  $k = 1, \dots, K$  with  $c_0 = -\sum_{k=1}^K c_k$ . Then since  $\psi$  is negative definite,

$$0 \geq \sum_{i=0}^K \sum_{j=0}^K c_i \psi(x_i, x_j) c_j$$

$$\begin{aligned} 0 \geq \sum_{i=0}^K \sum_{j=0}^K c_i \psi(x_i, x_j) c_j &= \sum_{i=1}^K \sum_{j=1}^K c_i \psi(x_i, x_j) c_j + c_0 \sum_{j=1}^K \psi(z, x_j) c_j + \\ & c_0 \sum_{i=1}^K c_i \psi(z, x_i) + c_0^2 \psi(z, z) \end{aligned}$$



# Positive and Negative

Proof.

$$\begin{aligned} \sum_{i=0}^K \sum_{j=0}^K c_i \psi(x_i, x_j) c_j &= \sum_{i=1}^K \sum_{j=1}^K c_i \psi(x_i, x_j) c_j - \sum_{i=1}^K c_i \sum_{j=1}^K \psi(z, x_j) c_j - \\ &\quad \sum_{i=1}^K \sum_{j=1}^K c_j c_i \psi(x_i, z) + \sum_{i=1}^K c_i \sum_{j=1}^K c_j \psi(z, z) \\ &= \sum_{i=1}^K \sum_{j=1}^K c_i (\psi(x_i, x_j) - \psi(x_i, z) - \psi(z, x_j) + \psi(z, z)) c_j \\ 0 &\geq - \sum_{i=1}^K \sum_{j=1}^K c_i \phi(x_i, x_j) c_j \\ 0 &\leq \sum_{i=1}^K \sum_{j=1}^K c_i \phi(x_i, x_j) c_j \end{aligned}$$

This make  $\phi$  positive definite. □

# Positive and Negative

Proof.

Suppose  $\phi$  is positive definite. Let  $c_1, \dots, c_K \in \mathbb{R}$  satisfy  $\sum_{k=1}^K c_k = 0$ . Then

$$\sum_{i=1}^K \sum_{j=1}^K c_i \phi(x_i, x_j) c_j \geq 0$$

Since  $\phi(x, y) = -\psi(x, y) + \psi(x, z) + \psi(z, y) - \psi(z, z)$

$$\begin{aligned} 0 &\leq \sum_{i=1}^K \sum_{j=1}^K c_i (-\psi(x_i, x_j) + \psi(x_i, z) + \psi(z, x_j) - \psi(z, z)) c_j \\ &\leq -\sum_{i=1}^K \sum_{j=1}^K c_i \psi(x_i, x_j) c_j + \sum_{i=1}^K c_i \psi(x_i, z) \sum_{j=1}^K c_j + \\ &\quad \sum_{i=1}^K c_i \sum_{j=1}^K \psi(z, x_j) c_j - \psi(z, z) \sum_{i=1}^K c_i \sum_{j=1}^K c_j = -\sum_{i=1}^K \sum_{j=1}^K c_i \psi(x_i, x_j) c_j \end{aligned}$$

Therefore  $\psi$  is negative definite. □

# Negative Definite From Positive Definite

## Proposition

Let  $K : \mathbb{R}^N \times \mathbb{R}^N \rightarrow \mathbb{R}$  be a positive definite symmetric function. Define  $\phi(x, y) = K(x, x) + K(y, y) - 2K(x, y)$ . Then  $\phi$  is negative definite.

## Proof.

Let  $x_1, \dots, x_M \in \mathbb{R}^N$  and  $c_1, \dots, c_M \in \mathbb{R}$  satisfying  $\sum_{m=1}^M c_m = 0$ . Then,

$$\begin{aligned} \sum_{i=1}^M \sum_{j=1}^M c_i \phi(x_i, x_j) c_j &= \sum_{i=1}^M \sum_{j=1}^M c_i (K(x_i, x_i) + K(x_j, x_j) - 2K(x_i, x_j)) c_j \\ &= \sum_{i=1}^M c_i K(x_i, x_i) \sum_{j=1}^M c_j + \sum_{j=1}^M c_j K(x_j, x_j) \sum_{i=1}^M c_i + \\ &\quad - 2 \sum_{i=1}^M \sum_{j=1}^M c_i K(x_i, x_j) c_j \leq 0 \end{aligned}$$



## Proposition

*Let  $K : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$  be a positive definite kernel satisfying  $K(x, y) \geq 0$ ,  $x, y \in \mathbb{R}^n$ . Then*

$$-\log K(x, y)$$

*is negative definite if and only if*

$$K(x, y)^t$$

*is positive definite for all  $t > 0$*

## Proposition

Let  $K : \mathbb{R}^N \times \mathbb{R}^N \rightarrow \mathbb{R}$  be a negative definite kernel that satisfies

$$K(x, y) > 0$$

Then,

$$\frac{1}{K(x, y)}$$

is a positive definite kernel.

# Other Positive Definite Kernels

- Laplacian:  $K(x, y) = \exp\left(-\frac{\|x-y\|}{\sigma}\right)$
- ANOVA:  $K(x, y) = \sum_{n=1}^N \exp(-\sigma(x^n - y^n)^2)^d$
- Rational Quadratic:  $K(x, y) = 1 - \frac{\|x-y\|^2}{\|x-y\|^2 + c}$
- Multiquadric:  $K(x, y) = \sqrt{\|x-y\|^2 + c^2}$
- Inverse Multiquadric:  $K(x, y) = \frac{1}{\sqrt{\|x-y\|^2 + c^2}}$
- Circular:  $K(x, y) = \frac{2}{\pi} \arccos\left(-\frac{\|x-y\|}{\sigma}\right) - \frac{2}{\pi} \frac{\|x-y\|}{\sigma} \sqrt{1 - \left(\frac{\|x-y\|}{\sigma}\right)^2}$   
if  $\|x-y\| < \sigma$ , zero otherwise
- Spherical:  $K(x, y) = 1 - \frac{3}{2} \frac{\|x-y\|}{\sigma} + \frac{1}{2} \left(\frac{\|x-y\|}{\sigma}\right)^3$  if  $\|x-z\| < \sigma$ , zero otherwise
- Wave Kernel:  $K(x, y) = \frac{\theta}{\|x-y\|} \sin \frac{\|x-y\|}{\theta}$
- Spline:  
$$K(x, y) = \prod_{n=1}^N \left(1 + x_n y_n + x_n y_n \min(x_n, y_n) - \frac{x_n + y_n}{2} \min(x_n, y_n)^2 + \frac{\min(x_n, y_n)^3}{3}\right)$$
- Cauchy:  $K(x, y) = \frac{1}{1 + \frac{\|x-y\|^2}{\sigma^2}}$
- Chi-Square:  $K(x, y) = 1 - \sum_{n=1}^N \frac{(x_n - y_n)^2}{\frac{1}{2}(x_n + y_n)}$
- Histogram:  $K(x, y) = \sum_{n=1}^N \min(x_n, y_n)$ , where  
 $x = (x_1, \dots, x_N)$  and  $y = (y_1, \dots, y_N)$